



Tombul *et al.* (2008) assessed 3-D computed tomography in 60 consecutive living kidney donors from 2002–2007.<sup>36</sup> Preoperative MDCT angiography detected 64 of the 67 renal arteries seen preoperatively in 60 renal units. Two undetected arteries had diameters less than 3 mm. The sensitivity of MDCT angiography was 95% for arteries and 93% for veins. The positive predictive value was 100% for arteries and veins. MDCT angiography was found to be less invasive and enabled rapid and accurate preoperative assessment of vascular anatomy in living kidney donors.

### Magnetic Resonance Imaging

Thirteen studies published from 1997 to 2006 compared operative findings with MRI angiographic findings.<sup>10,14,18,19,32,37–43</sup> The sensitivity in detecting accessory renal arteries ranged from 20%–100% (mean 80%). In studies with more than 100 participants, the mean sensitivity was 54%. This technique detects early branching with a mean sensitivity of 69%. It may miss fibromuscular dysplasia (incidence uncertain). Magnetic resonance angiography (MRA) source data is better than maximum intensity projection (MIP) data, which is better than virtual reality (VR) and shaded surface display (SSD) data.

Kok *et al.* (2008) evaluated the outcomes of vascular imaging and the clinical consequences of multiple arteries and veins.<sup>44</sup> Vascular anatomy at operation was compared with vascular anatomy as imaged by MRI or subtraction angiography. MRI failed to predict arterial anatomy in 23/220 compared with 3/101 after angiography. The authors concluded that both MRI and angiography provided suboptimal information on renal vascular anatomy.

Neville *et al.* (2008) prospectively compared MRA with selective renal angiography in patients from 53 renal units.<sup>45</sup> Selective renal angiography provided a sensitivity and specificity of 86% and 95%, respectively, and positive predictive value and negative predictive value of 75% and 97%, respectively. MRA had a sensitivity and specificity of 64% and 88%, respectively, and positive predictive value and negative predictive value of 58% and 90%, respectively. It was concluded that MRA could not replace standard renal angiography as the reference standard.

Monroy-Cuadros *et al.* (2008) retrospectively analysed the reliability of MRA compared with intra-operative findings in 66 patients.<sup>46</sup> In 8 cases, an accessory renal artery was found intra-operatively, 2 of which were incorrectly diagnosed as normal by MRA. The negative predictive value of MRA was 97%.

### SUMMARY OF THE EVIDENCE

CT evaluation is at least as good as CA and DSA in depicting detailed vascular anatomy of donor kidneys. Sixteen-slice CT machines may be superior to CA and DSA.

MRI may be slightly inferior to CT evaluation.

Both CT and MRI provide additional information about the renal parenchyma and urinary drainage of the kidneys. Both are less expensive to use than CA or DSA.

### WHAT DO THE OTHER GUIDELINES SAY?

**Kidney Disease Outcomes Quality Initiative:** No recommendation.

**UK Renal Association:** No recommendation.

**Canadian Society of Nephrology:** No recommendation.

**European Best Practice Guidelines:** The European Guidelines (ERA, EFPA) state that the use of contrast-enhanced ultrasound and conventional angiography with intravenous angiography being optional. The guideline was updated in 2000.

**International Guidelines:** No recommendation.

### SUGGESTIONS FOR FUTURE RESEARCH

Imaging modalities, especially MRI, are advancing rapidly in technological terms. This guideline is very likely to be out of date within 3 years and should be reviewed at the latest by 2011.

### CONFLICT OF INTEREST

Stephen Munn has no relevant financial affiliations that would cause a conflict of interest according to the conflict of interest statement set down by CARI.

### REFERENCES

- Gleeson MJ, Brennan RP, McMullin JP. Renal angiography in potential living related renal donors in Iraq. *Clin Radiol* 1988; **39**: 625–27.
- Sherwood T, Ruutu M, Chisholm GD. Renal angiography problems in live kidney donors. *Br J Radiol* 1978; **51**: 99–105.
- Figuerola K, Feuerstein D, Principe AL. A comparison of three-dimensional spiral computed tomography with renal angiography in the live-donor work-up. *J Transpl Coord* 1997; **7**: 195–98.
- Neymark E, LaBerge JM, Horose R *et al.* Arteriographic detection of renovascular disease in potential renal donors: incidence and effect on donor surgery. *Radiology* 2000; **214**: 755–60.
- Rubin GD, Alfrey EJ, Dake MD *et al.* Assessment of living renal donors with spiral CT. *Radiology* 1995; **195**: 457–62.
- Kaynan AM, Rozenblit AM, Figuerola KI *et al.* Use of spiral computerized tomography in lieu of angiography for preoperative assessment of living renal donors. *J Urol* 1999; **161**: 1769–75.
- Flechner SM, Sandler CM, Houston GK *et al.* 100 living-related kidney donor evaluations using digital subtraction angiography. *Transplantation* 1985 **40**: 675–78.
- Watarai Y, Kubo K, Hirano T *et al.* Intravenous digital subtraction angiography and helical computed tomography in evaluation of living renal donors. *Int J Urol* 2001; **8**: 417–22.
- Ayuso JR, Openheimer F, Ayuso C *et al.* Living donor kidney transplantation: helical CT evaluation of candidates. *Acta Urologica Espanolas* 2006; **30**: 145–151.
- Bakker J, Ligtenberg G, Beek FJ *et al.* Preoperative evaluation of living renal donors with gadolinium-enhanced magnetic resonance angiography. *Transplantation* 1999; **67**: 1167–72.
- Fink C, Hallscheidt PJ, Hosch WP *et al.* Preoperative evaluation of living renal donors: value of contrast-enhanced 3D magnetic resonance angiography and comparison of three rendering algorithms. *Eur Radiol* 2003; **13**: 794–801.

12. Hanninen EL, Denecke T, Stelter L *et al.* Preoperative evaluation of living kidney donors using multirow detector computed tomography: comparison with digital subtraction angiography and intraoperative findings. *Transplant Int* 2005; **18**: 1134–41.
13. Dachman AH, Newmark GM, Mitchell MT *et al.* Helical CT examination of potential kidney donors. *Am J Roentgenol* 1998; **171**: 193–200.
14. Bhatti AA, Chugtai A, Haslam P *et al.* Prospective study comparing three-dimensional computed tomography and magnetic resonance imaging for evaluating the renal vascular anatomy in potential living renal donors. *Br J Urol Int* 2005; **96**: 105–08.
15. Del Pozo JJ, Soria C, You-Cheng JW *et al.* Helical computerized tomography arteriography for evaluation of live renal donors undergoing laparoscopic nephrectomy. *J Urol* 1999; **162**: 31–34.
16. El-Diasty TA, Shokeir AA, El-Ghar ME *et al.* Contrast enhanced spiral computerized tomography in live kidney donors: a single session for anatomical and functional assessment. *J Urol* 2004; **171**: 31–34.
17. Feifer AH, Fong BC, Feldman L *et al.* Preoperative evaluation of laparoscopic living renal donors with computerized tomography and its effect on donor morbidity and graft function. *Can J Urol* 2005; **12**: 2713–21.
18. Halpern EJ, Mitchell DG, Wechsler RJ. Preoperative evaluation of living renal donors: comparison of CT angiography and MR angiography. *Radiology* 2000; **216**: 434–39.
19. El-Diasty TA, El-Ghar ME, Shokeir AA *et al.* Magnetic resonance imaging as a sole method for the morphological and functional evaluation of live kidney donors. *Br J Urol Int* 2005; **96**: 111–16.
20. Janoff DM, Davol P, Hazzard J *et al.* Computerized tomography with 3-dimensional reconstruction for the evaluation of renal size and arterial anatomy in the living kidney donor. *J Urol* 2004; **171**: 27–30.
21. Kapoor A, Kapoor A, Mahajan G *et al.* Multispiral computed tomographic angiography of renal arteries of live potential renal donors: a review of 118 cases. *Transplantation* 2004; **77**: 1535–39.
22. Kawamoto S, Montgomery RA, Lawler LP *et al.* Multidetector CT angiography for preoperative evaluation of living laparoscopic kidney donors. *Am J Radiol* 2003; **180**: 1633–38.
23. Kim JK, Park SY, Kim HJ *et al.* Living kidney donors: usefulness of multi-detector row CT for comprehensive evaluation. *Radiology* 2003; **229**: 869–76.
24. Lewis GR, Mulcahy K, Brook NR *et al.* A prospective study of the predictive power of spiral computed tomographic angiography for defining renal vascular anatomy before live-donor nephrectomy. *Br J Urol Int* 2004; **94**: 1077–81.
25. Lionel G, Sebben RA, Costello P *et al.* The use of spiral computed tomographic angiography for the assessment of living kidney donors. *Aust NZ J Surg* 1999; **69**: 217–19.
26. Namasivayam S, Small WC, Kalra MK *et al.* Multidetector-row CT angiography for preoperative evaluation of potential laparoscopic renal donors: how accurate are we? *Clinical Imaging* 2006; **30**: 120–26.
27. Patil UD, Ragavan A, Murthy NK *et al.* Helical CT angiography in evaluation of live kidney donors. *Nephrol Dial Transplant* 2001; **16**: 1900–04.
28. Pozniak MA, Balison DJ, Lee FT *et al.* CT angiography of potential renal transplant donors. *Radiographics* 1998; **18**: 565–87.
29. Rajamahanty S, Simon R, Edey M *et al.* Accuracy of three-dimensional CT angiography for preoperative vascular evaluation of laparoscopic living renal donors. *J Endourol* 2005; **19**: 339–41.
30. Raman SS, Pojchamarnwiputh S, Muangsombon K *et al.* Utility of 16-MDCT angiography for comprehensive preoperative vascular evaluation of laparoscopic renal donors. *Am J Radiol* 2006; **186**: 1630–38.
31. Ramirez-Bollas J, Hernandez-Dominguez M, Arenas-Osuna J *et al.* Preliminary report: usefulness of computed tomographic angiography in the protocol of a kidney donor. *Cirujia Cirujanos* 2003; **71**: 379–82.
32. Rankin JC, Jan W, Kofman CC. Noninvasive imaging of living related kidney donors: evaluation with CT angiography and gadolinium-enhanced MR angiography. *Am J Radiol* 2001; **177**: 349–55.
33. Rastogi N, Sahani DV, Blake MA *et al.* Evaluation of living renal donors: accuracy of three-dimensional 16-section CT. *Radiology* 2006; **240**: 136–44.
34. Sahani DV, Rastogi N, Greenfield AC *et al.* Multi-detector row CT in evaluation of 94 living renal donors by readers with varied experience. *Radiology* 2005; **235**: 905–10.
35. Johnston T, Reddy K, Mastrangelo M *et al.* Multiple renal arteries do not pose an impediment to the routine use of laparoscopic donor nephrectomy. *Clin Transplant* 2001; **15** (Suppl 6): 62–65.
36. Tombul ST, Aki FT, Gunay M *et al.* Preoperative evaluation of hilar vessel anatomy with 3-D computerized tomography in living kidney donors. *Transplant Proc* 2008; **40**: 47–9.
37. Ames SA, Krol M, Nettar K *et al.* Pre-donation assessment of kidneys by magnetic resonance angiography and venography: accuracy and impact on outcomes. *Am J Transplant* 2005; **5**: 1518–28.
38. Buzzas GR, Shield CF, Pay NT *et al.* Use of gadolinium-enhanced, ultrafast three-dimensional, spoiled gradient-echo magnetic resonance angiography in the preoperative evaluation of living renal allograft donors. *Transplantation* 1997; **64**: 1734–37.
39. Hodgson DJ, Wajanat J, Rankin S *et al.* Magnetic resonance renal angiography and venography: an analysis of 111 consecutive scans before donor nephrectomy. *Br J Urol Int* 2006; **97**: 584–86.
40. Israel GM, Lee VS, Edey M *et al.* Comprehensive MR imaging in the preoperative evaluation of living donor candidates for laparoscopic nephrectomy: initial experience. *Radiology* 2002; **225**: 427–32.
41. Low RN, Martinez AG, Steinberg SM *et al.* Potential renal transplant donors: evaluation with gadolinium-enhanced MR angiography and MR urography. *Radiology* 1998; **207**: 165–72.
42. Probst RL, Fernandez ED, Neff W *et al.* Evaluation of MR-angiography for pre-operative assessment of living kidney donors. *Clin Transplant* 2005; **19**: 522–26.
43. Steinborn M, Wintersperger BJ, Heck A *et al.* [Contrast enhanced MR angiography in the preoperative evaluation of living kidney donors]. *Rofo: Fortschritte auf dem Gebiete der Rontgenstrahlen und der Nuklearmedizin*. 1999; **171**: 313–18.
44. Kok NF, Dols LF, Hunink MG *et al.* Complex vascular anatomy in live kidney donation: imaging and consequences for clinical outcome. *Transplantation* 2008; **85**: 1760–65.
45. Neville C, House AA, Nguan CY *et al.* Prospective comparison of magnetic resonance angiography with selective renal angiography for living kidney donor assessment. *Urology* 2008; **71**: 385–9.
46. Monroy-Cuadros M, McLaughlin K, Salazar A *et al.* Assessment of live kidney donors by magnetic resonance angiography: reliability and impact on outcomes. *Clin Transplant* 2008; **22**: 29–34.